

Natura Beta

THE ANTI-STRESS PRODUCT OF CHOICE



1KG



6KG




20KG



Reduced sensitivity to environmental stress: temperature, water, and salt



Reduced cracking and increased assimilation of calcium.



Laevorotatory plant amino acids and readily available energy for the plant



Increased photosynthetic activity and cellular respiration: **BIOSTIMULATING ACTION**



Improvement of fruit quality characteristics (size, colour, texture, and shelf life)

NUTRITIONAL COMPOSITION	%
Total organic nitrogen (N)	5
Organic carbon (C) of biological origin	12
Plant-based organic matter	24
Glycine betaine	> 30
Laevorotatory plant amino acids	> 5

TECHNICAL SPECIFICATIONS	
S.W.	1.15
pH:	7

DOSAGE AND METHOD OF USE		Foliar application	Fertigation	
CROPS	PHENOLOGICAL STAGE	DOSE g/hl	DOSE kg/ha	NOTES
Stone and pome fruits	From flowering to fruit growth	300-600	15-20	applications every 7-15 days
Citrus trees	Flowering and fruit set	400-600	15-20	applications every 7-15 days
Actinidia	From flowering to fruit growth	400-600	15-20	applications every 7-15 days
Wine and table grapevines	Bunch closure and grape berry growth	300-600	15-20	applications every 7-15 days
Olive and hazel trees	Post-fruit set and fruit growth	400-600	15-20	applications every 7-15 days
Fruiting vegetables	Post-fruit set and fruit growth	300-600	8-15	applications every 7-15 days
Leafy vegetables	During the entire growing cycle	300-600	8-15	applications every 7-15 days
Root, bulb, or tuber vegetables	Starting from root, bulb, or tuber enlargement	300-600	8-15	applications every 7-15 days
Strawberries and small fruits	Post-fruit set and fruit growth	300-600	8-15	applications every 7-15 days
Tomatoes for industry	Post-fruit set and fruit growth	300-600	8-15	applications every 7-15 days
Industrial crops and legumes	Flowering and fruit set	300-600	/	applications every 7-15 days
Spring-summer cereals	Post-emergence and stem extension	300-600	/	applications every 7-15 days
Autumn-winter cereals	Tillering and flag leaf	300-600	/	applications every 7-15 days
Nursery, Ornamental Crops, and Floriculture	During the entire growing cycle	/	8-15	

- NOTES:**
- The dose is calculated based on a volume of spraying water of 8-10 hl/ha
 - Application is recommended in combination with MicroSync Ca, CaMa 104, or NitraCa918 to increase calcium absorption and reduce fruit diseases
 - Possible mixture with copper and sulphur on grapevines and olive trees



ORGANIC FARMING