

Natura Red

IMPECCABLE QUALITY AND DURABLE RIPENESS




6KG



25KG



Improvement of fruit quality characteristics (size, colour, texture, and shelf life)



Increased brix rating and uniform fruit veraison



Reduced over-ripening and increased shelf life



Macronutrients of vegetable origin that are highly available to plants

NUTRITIONAL COMPOSITION	%
Total organic nitrogen (N)	1.5
Organic potassium oxide (K ₂ O)	7
Organic carbon (C) of biological origin	15
Plant-based organic matter	30
Glycine betaine	1
Laevorotatory plant amino acids	> 7

TECHNICAL SPECIFICATIONS	
S.W.	1.25
pH:	7.5

DOSAGE AND METHOD OF USE		Foliar application	Fertigation	NOTES
CROPS	PHENOLOGICAL STAGE	DOSE g/hl	DOSE kg/ha	
Stone and pome fruits	Veraison and ripening	500-800	/	2-3 applications starting 30 days before harvest
Citrus trees	Veraison and ripening	800-1000	/	2-3 applications starting 30 days before harvest
Wine and table grapevines	Veraison and ripening	500-800	/	2-3 applications starting 30 days before harvest
Fruiting vegetables	Veraison and ripening	500-800	15-20	2-3 applications starting 30 days before harvest
Strawberries and small fruits	Veraison and ripening	500-800	15-20	2-3 applications starting 30 days before harvest
Tomatoes for industry	Veraison and ripening	500-800	/	2-3 applications starting when 40% of the berries are red

- NOTES:**
- The dose is calculated based on a volume of spraying water of 8-10 hl/ha
 - Apply during the coolest hours of the day, taking care to water the plants thoroughly
 - Do not apply in combination with alkaline-reacting pesticides or with products containing copper, sulphur, or mineral oils



ORGANIC FARMING