

Natura Sodi

SALINITY IS NO LONGER A PROBLEM



6KG



25KG



Improved chemical and physical properties of the soil and reduced soil fatigue



Reduced sensitivity to environmental stress: temperature, water, and salt



It reduces stress conditions caused by root asphyxia

NUTRITIONAL COMPOSITION

	%
Polyhydroxyphenylcarboxylic acids	40
Calcium oxide (CaO) soluble in water	7

TECHNICAL SPECIFICATIONS

S.W.	1.25
pH:	7

DOSAGE AND METHOD OF USE

CROPS	PHENOLOGICAL STAGE	Fertigation
		DOSE kg/ha
Stone and pome fruits	From flowering to fruit growth	5-10
Citrus trees	From flowering to fruit growth	5-10
Actinidia	From flowering to fruit growth	5-10
Wine and table grapevines	From flowering to grape berry growth	5-10
Olive and hazel trees	From flowering to fruit growth	5-10
Fruiting vegetables	Post-transplanting	1-1.5 kg/1000 sq.m.
Leafy vegetables	Post-transplanting	1-1.5 kg/1000 sq.m.
Root, bulb, or tuber vegetables	Post-transplanting	1-1.5 kg/1000 sq.m.
Strawberries and small fruits	Post-transplanting	5-10
Tomatoes for industry	Post-transplanting	5-10

- NOTES:**
- Do not mix with sulphates, phosphates, or alkaline-reacting products
 - Repeat with each fertigation in soils with high salinity